

The Canada-UK Deep Submillimetre Survey: First Submillimetre Images, the Source Counts, and Resolution of the Background

Stephen Eales¹, Simon Lilly², Walter Gear³, Loretta Dunne¹, J. Richard Bond⁴, Francois Hammer⁵, Olivier Le Fèvre⁶ and David Crampton⁷

ABSTRACT

We present the first results of a deep unbiased submillimetre survey carried out at 450 and 850 μm . We detected 12 sources at 850 μm , giving a surface density of sources with $S_{850\mu\text{m}} > 2.8 \text{ mJy}$ of $0.49 \pm 0.16 \text{ arcmin}^{-2}$. The sources constitute 20-30% of the background radiation at 850 μm and thus a significant fraction of the entire background radiation produced by stars. This implies, through the connection between metallicity and background radiation, that a significant fraction of all the stars that have ever been formed were formed in objects like those detected here. The combination of their large contribution to the background radiation and their extreme bolometric luminosities make these objects excellent candidates for being proto-ellipticals. Optical astronomers have recently shown that the UV -luminosity density of the universe increases by a factor of $\simeq 10$ between $z = 0$ and $z = 1 - 2$ and then decreases again at higher redshifts. Using the results of a parallel submillimetre survey of the local universe, we show that both the submillimetre source density and background can be explained if the submillimetre luminosity density evolves in a similar way to the UV -luminosity density. Thus, if these sources are ellipticals in the process of formation, they may be forming at relatively modest redshifts.

¹Department of Physics and Astronomy, Cardiff University, P.O. Box 913, Cardiff CF2 3YB, UK

²Department of Astronomy, University of Toronto, 60 St. George Street, Toronto, Ontario M5S 1A1, Canada

³Mullard Space Science Laboratory, University of London, Holmbury St. Mary, Dorking, Surrey RH5 6NT, UK

⁴CITA, University of Toronto, 60 St. George Street, Toronto, Ontario M5S 1A1, Canada

⁵DAEC, Observatoire de Paris-Meudon, 92195 Meudon, France

⁶Laboratoire d'Astronomie Spatiale, Traverse de Siphon, B.P.8 13376, Marseilles Cedex 12, France

⁷Dominion Astrophysical Observatory, 5071 W. Saanich Road, Victoria, BC, V8X 4M6, Canada

1. Introduction

The recent confirmation of a submillimetre background (Puget et al. 1996, Fixsen et al. 1998, Hauser et al. 1998) with approximately the same integrated intensity as the optical background (Pozzetti et al. 1998, Vogeley 1998) shows that to understand galaxy formation and evolution it is crucial to take into account that part of a galaxy's starlight that is absorbed by dust and re-radiated at longer wavelengths. The integrated background is the repository of all the stellar energy that has ever been emitted (with a $1+z$ weighting factor), and so the contribution of a particular class of galaxies to the background is a measure of its importance in the star-formation history of the universe. Determining the relative contribution to the background of different galaxy classes may be of particular importance for answering the question of how ellipticals and spiral bulges formed. The quantity of metals associated with nearby early-type galaxies (Edmunds and Phillipps 1997) implies that early-type galaxies should be responsible for about half the background (§4). Nevertheless, optical surveys have failed to find convincing evidence of proto-spheroids. Possible explanations of this failure are (a) that when the stars formed they were spread over a large volume of space before gradually coalescing to form galaxies (explaining the absence of discrete sources), (b) that the starbursts are hidden by dust. In both cases, these objects will still be contributing to the background; in the latter case they should be dominating the submillimetre background (Bond et al. 1986, Eales and Edmunds 1997).

The prospects for resolving the submillimetre background are good because of the commissioning of the SCUBA submillimetre array (Holland et al. 1998, Gear et al. 1998) on the James Clerk Maxwell Telescope. Last year a group carried out a submillimetre survey of two clusters, using the gravitational magnification by the clusters as an aid for finding distant submillimetre sources. They found four sources and concluded that optical surveys had significantly underestimated the number of star-forming galaxies in the distant universe (Smail et al. 1997, SIB, Blain et al. 1998). We and others (Barger et al. 1998, Hughes et al. 1998) have now started 'blind-field' surveys with SCUBA, which do not have the aid of gravitational amplification but do not suffer from the uncertainties of the cluster lens method. We are carrying out our survey in fields used for the Canada-France Redshift Survey (Lilly et al. 1995) and our aim is eventually to survey 200 arcmin² of the sky. In this paper we present the first results of a survey of an area of 22 arcmin² and analyse these using the results of a parallel survey we are carrying out of the submillimetre properties of the local universe (Dunne et al. 1998). A second paper (Lilly et al. 1998, hereafter Paper II) will discuss the optical counterparts to the distant submillimetre sources.

2. Observations and Data Reduction

We observed three fields simultaneously at 450 and 850 μm on 12 nights during January and March 1998, using the same observing procedure as SIB, with the exception that we chopped and nodded 30 arcsec EW. We divided our observations into integrations lasting about one hour, checking the telescope pointing before and after each integration. The optical depth during these nights varied between 0.57 and 1.65 at 450 μm and between 0.13 and 0.29 at 850 μm . For the first field we simply made repeated observations at the same position ($03^h 02^m 38.6^s 00^\circ 07' 40''$ [2000.0]), giving a total integration time of 36.096 ks. For the second field we made eight integrations (21.504 ks in total) at one position ($10^h 00^m 40.4^s 25^\circ 14' 20''$ [2000.0]) and four integrations (12.8 ks) at a second position ($10^h 00^m 38.4^s 25^\circ 14' 20''$ [2000.0]) 27 arcsec away from the first. For the final field we made observations in a mosaic pattern such that each point in the final map was constructed from data received at nine locations within the instrument field of view. Since the instrument is fixed in Naysmith coordinates, and thus the bolometers move on the sky, this ensures that a point on the final map contains data from a large fraction of the available bolometers, reducing the possibility of problems with individual bolometers generating spurious sources. We reduced our data in the standard way using the SURF package (Jenness 1997, Smail et al. 1997). The final maps are shown in Figs 1 & 2.

3. Source Extraction

Since any real object is likely to be unresolved and will appear on a map as a positive source with two negative sources 30 arcsec on either side, we used template analysis (Philips and Davies 1991) to look for sources of this kind on the 850- μm maps, using as a template a map of Uranus made with the same 30-arcsec chop. The negative emission is actually more complex than this because the chop direction is only precisely EW at the beginning of each integration, then remaining fixed at a constant azimuthal angle (this causes the slight rotation of the negative lobes visible in Fig. 2a), but this is a reasonable approximation. After trimming off the edge of each image, which is populated by noise artefacts due to incomplete sampling of the sky near the array's edge, we made both a map of the correlation coefficient (CC), a measure of how closely the emission resembles the template, and a map of the convolution of the image with the template (Fig. 2). The noise on the template-convolved images is 0.56, 0.76, and 0.92 mJy per beam for the 3^h , 10^h , and 14^h fields. We included a source in our catalogue only if both $S/N > 3$ on the convolved map and $CC > 0.4$, the latter criterion reducing the chance of noise spikes, which do not have the distinctive negative lobes, being included.

We checked the reality of our sources in two ways. First, we created the negative of each map and confirmed that no sources were found using the same selection criteria. Second, we generated Monte-Carlo simulations of the fields, replacing the data for each bolometer (which is sampled every second) with the output from a gaussian random-number generator with the same standard deviation as for that bolometer (measured separately for each one-hour integration). The simulations showed that spurious sources tended to be concentrated towards the edges of our fields as expected, and we used the simulations to define ‘reliable regions’ for each map (Figs 1 & 2). The sources listed in Table 1 are from these regions, and the simulations suggest that, on average, one of these could be spurious. Because of the possibility that the rotation of the chop direction could affect fluxes estimated by template-fitting, we also estimated fluxes by aperture photometry, finding no significant difference for the two methods.

Fig. 3 shows the integral source counts from our survey. An incompleteness analysis similar to that in SIB implies we are missing 0, 19, and 48% of the sources with fluxes of 6, 4, and 3 mJy respectively. Our source counts are consistent with those of the other small blank-field surveys (Barger et al. 1998, Hughes et al. 1998) and of SIB.

4. Discussion

The sources we have detected at $850\ \mu\text{m}$ account for 19% of the background emission at $850\mu\text{m}$ (Fig. 4), rising to 30% if a correction is made for incompleteness, and $>10\%$ of the background emission at $450\mu\text{m}$. Our four measurements and seven limits on the $450\mu\text{m}/850\mu\text{m}$ ratio are consistent with the average ratio being similar to that of the background emission, and thus consistent with the $850\mu\text{m}$ sources being representative of the background at shorter submillimetre wavelengths as well. On the assumption that the sub-mm objects detected in our survey make a negligible contribution to the optical background (Paper II), these objects account for $\sim 11\%$ of the total optical-near-IR plus submillimetre background (Dwek et al. 1998), rising to $\sim 17\%$ if the incompleteness correction is made.

It is possible that this level of background emission might be explained by AGN. If the black-hole masses measured in nearby galaxies by HST (Ford et al. 1998) are typical of galaxies in general, and if one makes the assumption that 10% of the mass accreting into a black hole is turned into energy, then the AGN background would be $\simeq 10\%$ of the background from stars. However, given the uncertainties in this calculation, we will henceforth make the more conservative assumption that we are observing emission from dust heated by stars.

The integrated background emission produced by a population at a redshift z is related to the smoothed out cosmic density of processed material $< \rho(Z + \Delta Y) >$ produced by these objects by $\int_0^\infty I_\nu d\nu = \frac{0.007 < \rho(Z + \Delta Y) > c^3}{4\pi(1+z)}$ (Pagel 1997). Since our sources make up a significant fraction of the background, they must represent a population which has been responsible for a significant fraction of all the star formation that has ever occurred. In Paper II we show that these sources are similar to the galaxies with huge far-infrared luminosities discovered with IRAS (Sanders and Mirabel 1996). This combination of high individual bolometric luminosities and a significant contribution to the background emission is exactly what one expects for one mode of proto-spheroid formation (§1), and it is hard to resist the conclusion that these are these long-sought objects (see also Barger et al. 1998, Hughes et al. 1998). Note that, with their huge bolometric luminosities and implied star-formation rates, it has long been suspected that ultra-luminous IRAS galaxies might be ellipticals in the process of formation (Wright et al. 1990). What has been lacking is direct evidence that objects like this at high redshift are responsible for a significant fraction of the star formation in the Universe.

An obvious question to ask is whether these objects are at $z \sim 1 - 2$, as are the galaxies that dominate the optical background (Pozzetti et al. 1998), or whether they are at higher redshifts, as one might suspect from the ages of the stellar populations in nearby ellipticals (Bower et al. 1992). We will address this question by making the null hypothesis that the submillimetre luminosity density evolves in the same way as the UV luminosity density (Madau et al. 1998) and predicting the submillimetre source counts and background. If the observed counts and background exceed our predictions, this might be evidence for a population of galaxies at $z \gg 1$. In Paper II we will examine this question using the optical identifications.

The biggest problem in modelling the submillimetre source counts and background has been the lack of information about the submillimetre properties of nearby galaxies. Without this information, one is forced to start from a local IRAS 60- μm luminosity function and make an assumption about the average far-infrared/submillimetre SED of galaxies, an extrapolation of a factor of over 10 in wavelength (Eales and Edmunds 1996). To overcome this problem, we have been carrying out a parallel submillimetre survey of the local universe (Dunne et al. 1998). By measuring submillimetre fluxes for galaxies drawn from statistically-complete samples of IRAS and optically-selected galaxies, we can both determine the SED's for different types of galaxy and (using accessible-volume techniques—Avni and Bahcall 1980) the local submillimetre luminosity function. As yet our local 850- μm luminosity function is only well constrained at high submillimetre luminosities, which is acceptable for making source-count predictions, since it is only the high-luminosity sources which contribute at the flux densities of interest, but not for

estimating the background. Fig. 3 shows our source-count predictions. We implemented our null hypothesis in two ways: (i) by having the amplitude of the local luminosity function evolve with the same form as the luminosity density; (ii) by having the luminosities of the sources evolve with this form. Fig. 3 shows that with luminosity evolution, but not with space-density evolution, the predicted counts match the observed counts fairly well. If one assumes no evolution at all, the observed source counts exceed the predicted counts by a large factor, as noted by Smail et al. 1997.

For modelling the background we are thrown back on the IRAS 60- μm luminosity function (Lawrence et al. 1986), but we do have the advantage over previous studies of empirical results on the far-IR-submillimetre SED's of galaxies. The 60, 100, and 850- μm fluxes of IRAS galaxies are well fitted by a single-temperature dust spectrum with a dust-emissivity index of 1 and $T_d = 7.5 \log_{10} L_{FIR} - 42.5$ over the range $10.0 < \log_{10} L_{FIR} < 12.5$, L_{FIR} being defined as in Lawrence et al. (Dunne et al. 1998). We do not claim that the dust-emissivity index is 1 and that there is dust of only one temperature in these galaxies, merely that this is a good empirical fit to the fluxes. Fig. 4 shows predictions using the IRAS luminosity function and this assumption about galaxy SED's. The measured background is far in excess of the no-evolution model, but the model in which the submillimetre luminosity density evolves in the same way as the UV luminosity density matches the spectral shape and amplitude of the background remarkably well.

Our conclusion that the null hypothesis of evolution like that in the optical waveband is acceptable (further supported by the analysis in Paper II) should be contrasted with recent claims to the contrary (Blain et al. 1998, Hughes et al. 1998). The difference is not in the data, but in the assumptions made about the local SED's. Blain et al., for example, assume $T_d = 38$ K and $n = 1.5$. When compared with our assumptions about SED's, this makes a factor of $\simeq 9$ difference in intensity at 850 μm .

The level of the background can also be used to argue against a major population of dust-enshrouded proto-ellipticals at $z > 5$. Let us assume that all the stars in this hypothetical population form instantaneously at a redshift z_F and that the SED from one of these objects is the same as that of the archetypical local IRAS galaxy Arp 220 (the precise SED is not important). This is enough to define the *shape* of the background. The maximum amplitude of this background is then set by the observed COBE background (Fig. 4). We then subtract this maximal elliptical background from the observed background to get a remnant background, which we assume is from the spiral/irregular population. If we then assume that late-type galaxies go through their period of maximum star formation at $z \sim 1$, we can then use the equation above and the ratio of the two backgrounds to predict the relative contributions to the present-day metal abundance of the two populations. If

$z_F = 5$ the dusty ellipticals can produce $<38\%$ of the metals of the other population and if $z_F = 10$ this falls to $<8\%$. Observations suggest, however, that the ratio of the global metal abundance associated with ellipticals to that associated with spirals/irregulars lies between 1 and 6 (Pagel 1997, Edmunds and Phillipps 1997). This discrepancy again suggests that ellipticals form at relatively modest redshifts.

We are grateful to the many people that have contributed to the development of SCUBA and to the JAC staff for their excellent support during this program, in particular Wayne Holland for help in a multitude of areas. We also thank Rob Ivison for some useful advice. Loretta Dunne is supported by a PPARC studentship. Research by Dick Bond and Simon Lilly is supported by the Natural Sciences and Engineering Research Council of Canada and by the Canadian Institute of Advanced Research. The JCMT is operated by the Joint Astronomy Center on behalf of the UK Particle Physics and Astronomy Research Council, the Netherlands Organization for Scientific Research and the Canadian National Research Council.

Table 1. Source Catalogue.

Name	RA (2000.0)	Dec (2000.0)	S/N	CC	S _{850μm} (mJy)	S _{450μm} (mJy)
CFRS03A	03 02 36.27	00 08 13	6.1	0.61	3.4 $\pm$ 0.6	17 $\pm$ 7
CFRS10A	10 00 38.19	25 14 50	4.2	...	4.8 $\pm$ 1.0	23 $\pm$ 9
CFRS10B	10 00 37.10	25 14 58	5.0	...	4.7 $\pm$ 1.0	21 $\pm$ 8
CFRS10C	10 00 37.23	25 15 12	3.8	...	2.8 $\pm$ 1.0	<22
CFRS10D	10 00 36.97	25 14 42	3.6	...	3.9 $\pm$ 1.0	<22
CFRS14A	14 17 40.3	52 29 08	8.8	0.78	8.8 $\pm$ 1.1	<31
CFRS14B	14 17 51.8	52 30 29	6.3	0.64	5.2 $\pm$ 0.9	<31
CFRS14C	14 17 33.6	52 30 49	3.4	0.46	3.8 $\pm$ 1.1	<31
CFRS14D	14 18 02.5	52 30 53	3.6	0.47	3.2 $\pm$ 0.9	<31
CFRS14E	14 18 02.6	52 30 17	3.6	0.54	3.5 $\pm$ 1.0	<31
CFRS14F	14 17 42.2	52 30 27	3.1	0.43	2.7 $\pm$ 0.9	20 $\pm$ 11
CFRS14G	14 17 29.7	52 31 15	3.7	0.45	5.7 $\pm$ 1.6	...

Note. — Col. 1: name of source. Cols 2 & 3: position (2000.0). The positions were measured from the 850- μ m maps, except for those of CFRS03A, CFRS10A, and CFRS10B, which were measured from the 450- μ m maps. Col. 4: Signal-to-noise. This is the peak signal on the map produced by convolving the raw map with the template (see text) divided by the noise on that map, except in the cases of the 10-hour sources, where the signal-to-noise was measured from the raw map. Col. 5: the correlation coefficient. The values for the 10^h sources are missing because template-fitting was not used for this field. Col. 6: flux at 850 μ m. These were obtained by template-fitting, except those for the 10-hour sources were obtained by aperture photometry, using an aperture of diameter 20 arcsec. The errors include a 5% calibration error. Col. 7: fluxes at 450 μ m. These were obtained by aperture photometry, using an aperture of diameter 12 arcsec, except that the flux for CFRS03A was obtained by template-fitting. The errors include a calibration error of 20%. The limits are 3 σ upper limits. There is no measurement for CFRS14G because the slightly smaller field-of-view of SCUBA at 450 μ m means that at this wavelength it falls in a noisy edge region.

REFERENCES

- Avni, Y. & Bahcall, J.N. 1980, ApJ, 235, 694.
- Barger, A.J. et al., Nature, in press.
- Blain, A.W., Smail, I., Ivison, R.J. & Kneib, J-P. 1998, MNRAS, submitted (astro-ph 9806063).
- Bond, J.R., Carr, B.J. & Hogan, C.J. 1986, ApJ, 306, 428.
- Bower, R.G., Lucey, J.R. & Ellis, R.S. 1992, MNRAS, 254, 601.
- Dunne, L. et al., in preparation.
- Dwek, E. et al. 1998, ApJ, in press.
- Eales, S.A. & Edmunds, M.G. 1996, 280, 1167.
- Eales, S.A. & Edmunds, M.G. 1997, MNRAS, 286, 732.
- Edmunds, M.G. & Phillipps, S. 1997, MNRAS, 292, 733.
- Fixsen, D.J. et al. 1998, ApJ, in press.
- Ford, H.C., Tsvetanov, Z.I., Ferrarese, L. & Jaffe, W. 1998, The Central Regions of the Galaxy and Galaxies, IAU 184, in press (astro-ph 9711299).
- Gear, W. et al., in preparation.
- Hauser, M.G. et al. 1998, ApJ, in press.
- Holland, W.S. et al. 1998, Advanced Technology MMW, Radio and Terahertz Telescopes, ed. Phillips, T., Proc SPIE 3357, 1998.
- Hughes, D.H. et al. 1998, Nature, in press.
- Jeness, T. 1997, SURF - SCUBA user reduction facility. *Starlink User Note 216.1*.
- Lawrence, A., Walker, D., Rowan-Robinson, M., Leech, K.J. & Penston, M.V. 1986, MNRAS, 219, 687.
- Lilly, S., Eales, S.A., Gear, W., Hammer, F., Le Fèvre, O., Crampton, D., Bond, J.R. & Dunne, L., in preparation.
- Lilly, S., Le Fèvre, O., Crampton, D., Hammer, F. & Tresse, L. 1995, ApJ, 455, 50.

- Madau, P., Della Valle, M. & Panagia, N. 1998, MNRAS, 279, L17.
- Pagel, B. 1997, Nucleosynthesis and Chemical Evolution of Galaxies (CUP).
- Phillips, S. & Davies, J. 1991, MNRAS, 251, 105.
- Pozzetti, L., Madau, P., Zamorani, G., Ferguson, H.C. & Bruzual, G.A. 1998, MNRAS, submitted (astro-ph 9803144).
- Puget, J-L. et al. 1996, A&A, 308, L5.
- Sanders, D.B. & Mirabel, I.F. 1996, ARA&A, 74, 749.
- Smail, I., Ivison, R.J. and Blain, A.W. 1997, ApJ, 490, L5.
- Vogele, M.S. 1998, ApJ, submitted (astro-ph 9711209).
- Wright, G.S., James, P.A., Joseph, R.D. & McLean, I.S. 1990, Nature, 344, 417.

Fig. 1.— (Plate 1) Images at $850\ \mu\text{m}$ of the 3^h (1a) and the 10^h (1b) fields. The beam of the telescope has a FWHM of 14 arcsec at $850\ \mu\text{m}$. The 3^h image has been convolved with a 14-arcsec gaussian (FWHM), the 10^h image with an 8-arcsec gaussian. The contour lines are drawn at $\pm 2, 3, 4, 5 \dots \sigma$. The dashed line shows the region which is believed to be free of edge artefacts and to have reasonably uniform noise properties. This region is $\simeq 110$ arcsec across for the 3^h field and $\simeq 120$ across (in the vertical direction) for the 10^h field.

Fig. 2.— (Plate 2) Images at $850\ \mu\text{m}$ of the 14^h field. The upper image (2a) has been convolved with a 14-arcsec gaussian, the lower image (2b) has been convolved with the source template (see text). Both images have been divided by the square root of a variance array produced from a Monte-Carlo simulation of the field, which corrects for uneven noise across the images. The contour lines are drawn at $\pm 2, 3, 4, 5 \dots \sigma$. As the convolution with the template uses information from both the positive and negative ‘lobes’, there are more prominent sources in 2b than in 2a, although one has to beware spurious sources being created 60 arcsec EW from a bright source (e.g. source Z). The dashed line in the lower image shows the region which is believed to be free of edge artefacts. This is $\simeq 390$ arcsec across in the horizontal direction.

Fig. 3.— Integral source counts at $850\ \mu\text{m}$ from our survey (dots) and other surveys: triangle (Hughes et al. 1998), circle (Barger et al. 1998) and cross (Smail et al. 1997). The errors on our counts are Poisson errors and are not independent. No correction has been to any of the counts for incompleteness, although the incompleteness for the other two blind-field surveys is claimed to be low. The predictions were made as described in the test. The key is as follows: no evolution, $\Omega_0 = 1$ —squares; density evolution, $\Omega_0 = 0$ —dots in circles; density evolution, $\Omega_0 = 1$ —circles; luminosity evolution, $\Omega_0 = 0$ —stars; luminosity evolution, $\Omega_0 = 1$ —triangles.

Fig. 4.— The dashed line and the points show, respectively, the submillimetre background estimated from the results of the FIRAS (Fixsen et al. 1998) and DIRBE (Hauser et al. 1998) experiments on COBE. The two lower limits show the limits on the background produced by simply adding up the number of sources that we observe. The solid line and the dotted lines show backgrounds predicted using an IRAS luminosity function and information on the SED’s of galaxies from our nearby galaxy survey. The solid line is for no evolution, the dotted line for a model in which the submillimetre luminosity density evolves like the UV luminosity density. The two other lines are the backgrounds predicted for a model in which all the stars in ellipticals form instantaneously at a redshift z_F , with the stars being hidden by dust. The predicted backgrounds have been normalized so that they are the maximum backgrounds consistent with the FIRAS measurement. The higher curve is for $z_F = 5$, the lower for $z_F = 10$.











